AWARD NUMBER: W81XWH-15-1-0211

TITLE: Cognitive and Neural Correlates of Aging in Autism Spectrum Disorder

PRINCIPAL INVESTIGATOR: Leslie C. Baxter, PhD

CONTRACTING ORGANIZATION: St. Joseph's Hospital and Medical Center

Phoenix, AZ 85013

REPORT DATE: July 2017

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;

Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

		Forms America d
REPORT DO	CUMENTATION PAGE	Form Approved
		OMB No. 0704-0188 nstructions, searching existing data sources, gathering and maintaining the
data needed, and completing and reviewing this collection this burden to Department of Defense, Washington Headqu	of information. Send comments regarding this burden estimate or any othe lariters Services, Directorate for Information Operations and Reports (0704- any other provision of law, no person shall be subject to any penalty for fail	r aspect of this collection of information, including suggestions for reducing
1. REPORT DATE	2. REPORT TYPE	3. DATES COVERED
July 2017	Annual	1 Jul 2016 - 30 Jun 2017
4. TITLE AND SUBTITLE		5a. CONTRACT NUMBER
"Cognitive and Neural Correlates of Aging in Autism Spectrum Disorder"		5b. GRANT NUMBER W81XWH-15-1-0211
		5c. PROGRAM ELEMENT NUMBER
6. AUTHOR(S)		5d. PROJECT NUMBER
Leslie C. Baxter, PhD; Brittany E	Blair Braden, PhD	
leslie.baxter@dignityhealth.org		E TAGICAUMPED
		5e. TASK NUMBER
		5f. WORK UNIT NUMBER
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)	8. PERFORMING ORGANIZATION REPORT NUMBER
St. Joseph's Hospital and Medical (350 West Thomas Road Phoenix, AZ 85013	Center	
9. SPONSORING / MONITORING AGENCY U.S. Army Medical Research and M		10. SPONSOR/MONITOR'S ACRONYM(S)
Fort Detrick, Maryland 21702-5012		11. SPONSOR/MONITOR'S REPORT NUMBER(S)
12. DISTRIBUTION / AVAILABILITY STAT	EMENT	<u>'</u>
Approved for Public Release; Distri		
13. SUPPLEMENTARY NOTES		

14. ABSTRACT

Purpose: As the first diagnosed ASD individuals are now reaching old age, it is imperative that we understand the impact of aging on individuals with ASD. We developed a model predicting greater executive dysfunction and frontal lobe susceptibility in ASD beyond normal aging. Scope: This study, which is a collaborative study of the Southwest Autism Research and Resource Center and the Barrow Neurological Institute, produces comprehensive cognitive, behavioral, and neuroimaging data on a set of well-characterized older ASD individuals who can be used as a reference for clinical diagnosis, therapeutics, and care plans. The **scope** of the second year was to continue with recruitment and data collection, and produce results from cross-sectional analyses of the initial assessment. **Results and significance:** We published our first in a series of studies from the initial cross-sectional data analysis in a high impact Autism journal (Autism Research). We found that the older ASD group performed significantly worse in executive functioning and showed decreased white matter connectivity in the frontal and temporal lobes. We also submitted resting state data regarding frontal connectivity group differences to the Society for Neuroscience meeting and Dr. Braden was selected to chair a nanosymposium on Autism: Physiology and Behavior. We have two additional data analyses on other brainbehavior relationships that are being prepared for publication, and we continue to collect data from new recruits and from returning participants.

15. SUBJECT TERMS

Autism, aging, functional MRI, fMRI, neuroimaging, executive functioning, memory, cognition, cortical thickness, connectivity, white matter, sparse Bayesian networks, machine learning

16. SECURITY CLASSIFICATION OF:		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON USAMRMC	
a. REPORT	b. ABSTRACT	c. THIS PAGE	Unclassified	18	19b. TELEPHONE NUMBER (include area code)
Unclassified	Unclassified	Unclassified	0.10.00000		

Standard Form 298 (Rev. 8-98) Prescribed by ANSI Std. Z39.18

Table of Contents

	<u>. I</u>	age
1. I	ntroduction	5
2. K	Keywords	5
3. A	Accomplishments	5
4. Iı	mpact	13
5. C	hanges/Problems	14
6. P	Products	15
7. P	articipants & Other Collaborating Organizations	17
8. S	pecial Reporting Requirements	
9. A	Appendices	

1. INTRODUCTION:

As the first diagnosed Autism Spectrum Disorder (ASD) individuals are now reaching old age, it is imperative that we understand the impact of aging on individuals with ASD. Given the striking parallels in ASD of deficits in executive function, subserved by the frontal lobe, and that the frontal lobe is susceptible to normal age-related changes, we combine neuroimaging, cognitive assessments and behavioral measures to examine aging in ASD compared to Typically Developed (TD) adults. We hypothesize that individuals with ASD will have an exacerbation of deficits beyond normal aging, as evidenced by significantly lower scores on tests affected by aging (e.g., executive) along with neuroanatomical markers of dysfunction, and relative preservation of function subserved by more posterior brain regions (memory and local detail processing). Our objective is to produce comprehensive cognitive, behavioral, and neuroimaging data on a group of well-characterized older individuals with ASD who can be used as a reference for clinical diagnosis, therapeutics, and care plans. To achieve this goal, our three year project involves longitudinal assessment of aging (40–60 y.o.) ASD individuals versus age-matched TD. In addition to commonly used statistical methods, we will use innovative machine learning and sparse Bayesian networks to combine structure, function, cognition, and symptom profiles to specifically address contributions to accelerated aging in ASD.

2. **KEYWORDS:** Autism, aging, functional MRI, fMRI, neuroimaging, executive functioning, cognition, memory, white matter, cortical thickness, connectivity

3. **ACCOMPLISHMENTS:**

What were the major goals of the project?

The major goals for the project in the second year of funding were:

- A) To continue obtain cognitive and MRI data from ASD and control participants (Major Task 2 and Subtask 3).
- B) Analyze and synthesize data to address all specific aims (Major Task 3, Milestones 2).

What was accomplished under these goals?

A) Obtaining cognitive and MRI data from ASD and control participants: We continue to collect new participants to obtain our target enrollment of 70 participants (35 per group).

At the time of this writing, we have all procedures for Time 1 for 46 of 70 participants. We have also performed the second time point for the initial cohort of participants who reached their 2 year interval time point. We have been able to complete all the procedures for 21 of 24 participants. Two of the ASD participants list above did not receive MRI scans because they were uncomfortable in the scanner, but we obtained cognitive and other data. The rest of the participants tolerated the scanning procedure well; none of those participants have required any alteration of the MRI or cognitive protocols. In our original Statement of Work, recruitment was scheduled for 3-18 months. Recruitment lags somewhat behind our anticipated schedule, although we continue to receive a small but regular flow of potential recruits. As soon as our new paper is available for viewing (see below), we plan to have increased media exposure, which tends to increase self-referrals.

C) Analyze and synthesize data to address all specific aims (Major Task 3, Milestones 2). We have published the first set of results from our initial cross-sectional analyses in Autism Research, the official journal of the International Society for Autism Research (Impact factor: 3.765; ranked 5/51 for Behavioral Sciences and 9/70 for Psychology Developmental journals). We are currently finishing data analyses and manuscript preparation for two more cross-sectional studies examining other cognitive functions presented below. We have extended our analyses to examine functional connectivity using resting state scans (re-FC) to examine the relationship with symptoms. We have also applied for and received funding to expand our study to include aging women with and without ASD who will be used with the current data to investigate gender differences in brain, cognition and symptoms. Details of significant data results are detailed below.

Significant results from cross-sectional analyses:

"Executive Function and Functional and Structural Brain Differences in Middle-Age Adults with Autism Spectrum Disorder" B. Blair Braden, PhD¹, Christopher J. Smith, PhD², Amiee Thompson³, Tyler K. Glaspy, BS³, Emily Wood³, Divya Vatsa³, Angela Abbott³, Samuel C. McGee³, and Leslie C. Baxter, PhD³. Accepted for publication: **Autism Research, DOI10.1002/aur.1842**. ¹Department of Speech and Hearing Science, Arizona State University, 976 S Forest Mall, Tempe, AZ 85281


²Southwest Autism Research & Resource Center, 2225 N 16th Street, Phoenix, AZ 85006 ³Department of Neuroimaging Research, Barrow Neurological Institute, St. Joseph's Hospital and Medical Center, 350 W Thomas Rd, Phoenix, AZ 85013

Our published report included analysis of cognitive and fMRI data from the first 16 ASD and 17 cognitively normal ("neurotypical" or NT) controls. All ASD diagnoses were confirmed by the trained staff at Southwest Autism Resource and Research Center (SARRC) based on the

Autism Diagnostic Observation Schedule. All participants were given the Kaufman Brief Intellectual Test to provide an estimate of intelligence to ensure that all participants were above the IQ= 80 cutoff and to assure general equivalence of IQ between groups. Groups were well-matched according to age, IQ, and education. Groups performed similarly on tests of delayed verbal memory (RAVLT) and detailed visual search (Group Embedded Figures Task), but the ASD group made significantly more errors on the executive function task, which requires monitoring, attention, mental flexibility and concept formation (Wisconsin Card Sorting Test (WCST); Table 1). A significant group difference on WCST errors remained after including demographic variables (age, IQ, or education) as covariates (all p<0.05). On fMRI tasks, performance was above 80% on all tasks.

Using independent component analysis (ICA), we evaluated functionally connected neural network activity during the n-back task. The n-back task is a common fMRI task used to interrogate the integrity of networks involved in working memory. The task requires online monitoring and attention to a series of letters. The ASD participants' reaction time was slower on the 2-back condition. We demonstrated older adults with ASD have weaknesses on an executive function task emphasizing flexibility, working memory, and inhibition (WCST) and are less likely to engage a CSTC network when switching between low-load and high-load working memory conditions. The beta weights were more positive for the 2-back condition [F(1,28) =11.182; p = 0.002] and more negative for the 0-back condition [F(1,28) = 17.471; p < 0.001] for the NT group. Striato-thalamo-cortical networks are known to be involved in working memory, impulse control and other aspects of executive functioning (Miller and Cohen, 2001). Recently, Gordon et al. (2015) demonstrated that differences in functional connectivity between regions of this subcortical-frontal network were associated with variability in normal controls. Specifically, lack of engagement of this network resulted in weak performance on the n-back task and increased impulsivity scores within a group of healthy controls. The findings from our group differences in engagement of this network along with the greater number of errors produced by the ASD group on the WCST, which requires error monitoring, updating and flexible thinking, suggests that this is a relative weakness within our ASD group, detectible within a small group. We will begin collecting our second longitudinal data point for those already enrolled in the study to determine whether aging exacerbates these differences.

Figure 1. Cortico-striatal-thalamo-cortical network differences between ASD and Control groups



White matter:

The ASD group showed reduced white matter integrity (voxel-wise differences in Fractional Anisotropy; FA) in the genu of the corpus callosum and bilaterally in the fimbria of the hippocampi. There were no areas of greater FA in the ASD cohort. No group differences in Mean Diffusivity, another measure of white matter integrity survived correction for multiple comparisons, but consistent with the spatial location of FA differences, two clusters in the bilateral anterior corona radiata (projections from of the genu of the corpus callosum) showed increased MD in the ASD group at p=0.001 uncorrected (not shown).

"Autism and age-related network differences in verbal fluency in adult autism"

Baxter LC¹, Wood E¹, Singh K¹, Smith CJ², Braden BB³.

Language functioning is variable in autism, ranging from nonverbalism in more severely affected individuals, to pragmatic aspects of language comprehension and mild problems with language processing in high functioning individuals. Using a commonly used fMRI fluency task, we determined the effects of ASD and aging on language networks.




Figure 2. Fluency task results across all groups, showing robust activation in left inferior frontal lobe ("Broca's area" and also right hemisphere cerebellar activation.

Fluency tasks are complex tasks that combine language and executive functions since they require the individual to engage associative processes and strategies to search and produce words according to a given demand (e.g., words with the same first letter). In our study, participants perform a "fluency" fMRI task, which is based on the Controlled Oral Word Association Test (COWAT), a task often used to assess expressive language abilities. This is a commonly-used fluency paradigm, requiring the participant to silently generate as many words they can to a designated first letter. The task has blocks of 16 seconds, during which the participant thinks of words to two letters (i.e., 8 seconds per letter) followed by a baseline block during which the word

"Relax" is presented and the participant is instructed to not talk to themselves, and encouraged to "think of a blank blackboard". The participant presses a button with their right forefinger for each word they generate to estimate the rate of word generation. We analyzed cross-sectional data from middle age (40-60) and young adult (18-25) high-functioning ASD and age-match TDs. There were no significant group differences for fMRI task performance, as measured by mean number of words produced on the task (about 50 words, p = .66). We first used ICA (described above) for each of the 4 groups to visualize any network differences across groups. We rejected any components indicating movement or the motor network associated with finger


¹Department of Neuroimaging Research, Barrow Neurological Institute, Phoenix, AZ


²Southwest Autism Research & Resource Center, Phoenix, AZ


³Speech and Hearing Sciences, Arizona State University, Tempe, AZ

movement. We found several components that significantly correlated with the task. All groups produced a network involving left inferior frontal cortex (LIFC), in the general area often termed "Broca's area" that is critical for expressive language. Other networks were found that included the right cerebellum, anterior cingulate, left hippocampus, and the precuneus/posterior cingulate region reflecting engagement of this network during the baseline condition.

Using SPM, we extracted values from any nodes found from networks significantly engaged during any of each of the 4 group ICA analyses in order to test for group, age, and group-by-age interactions. Our results indicated that not all of the groups utilized these regions to the same extent when performing the fluency task, despite all groups having very similar task performance. Specifically, some regions contributing to this fluency task showed clear aging effects, while others showed differences between the ASD group and the TD groups (Figure 2). For example, the left hippocampus was far less engaged in the fluency task for both older ASD and TD individuals; in contrast, both older and younger adult ASD groups showed less cerebellar activation compared to the TD groups. Interestingly, the older ASD group shows a level of engagement that is similar to young controls, possibly suggesting a compensatory mechanism.










Figure 3. Regional BOLD activity during a fluency task. Both ASD groups show decreased right cerebellar activity while both older groups engage the hippocampus to a much lesser extent than their younger cohorts. The ASD groups show less engagement of the precuneus region that is part of the Default Mode Network during the baseline rest condition. The older ASD group shows a relatively high degree of engagement of Broca's area, possibly as a compensation for weaknesses in other regions. All y axes are contrast estimates obtained from whole brain analysis in SPM(12).

"Motor system integrity in older adults with autism spectrum disorder"

Deatherage BR¹, Braden BB², Smith CJ³, McBeath MM⁴, Singh K¹, Thompson AM¹, Wood EG¹, Vasta D¹, McGee S¹, Baxter LC¹

We are finalizing a manuscript that in part includes an honor's project by ASU undergraduate student Brandon Deatherage. Gait disturbance, clumsiness, and other mild movement problems are often observed in children with autism spectrum disorder (ASD) (Maurer, 1982). As the brain ages, these symptoms may persist into late adulthood in those diagnosed with ASD. This study focused on middle-age adults with ASD to study motor behavior and underlying brain integrity in older individuals. Using a finger tapping task, motor performance was measured in a cross-sectional study comparing middle-aged adults with ASD and age-, education- and IQ-matched typically developing (TD) controls. We hypothesized that older adults with ASD would show poorer motor performance (slower finger tapping speed). We also hypothesized that underlying brain differences, measured using MRI, in regions associated with motor function including the primary motor cortex, basal ganglia, and cerebellum, as well as the white matter connecting tracts would exist between groups and may explain the proposed disparity in motor performance.

Data from the study show that older ASD participants have greater variability in right hand finger tapping than the TD group, although there was not a significant difference in mean group values. On a brain level, the left motor cortical region was the only gray matter region showing a significant effect, with thinning occurring in the ASD group. In contrast, group differences were found on several measures of white matter including the cortical spinal tract, posterior limb of the internal capsule, retrolenticular internal capsule and middle cerebellar peduncle. Brain-behavior correlations showed that dominant finger tapping speed, which showed greater variability among the ASD group, correlated with left hemisphere white matter integrity measures of the left corticospinal tract and posterior limb and retrolenticular segment of the internal capsule and cerebellar white matter.

No significant differences were observed between groups in finger tapping speed but the dominant, right-hand tapping showed greater variability in the ASD cohort. Decreased white matter integrity was observed in the ASD group for many regions associated with motor functioning. Given that white matter integrity, not gray matter thickness, correlated with right

¹Department of Neuroimaging Research, Barrow Neurological Institute, Phoenix, AZ

²Speech and Hearing Sciences, Arizona State University, Tempe, AZ

³Southwest Autism Research & Resource Center, Phoenix, AZ

⁴Department of Psychology, Arizona State University, Tempe, AZ

hand tapping in the ASD suggests that white matter differences may account for the heterogeneity and intermittent findings of motor dysfunction in ASD groups.

Other related accomplishments:

Received funding to include a cohort of older women for longitudinal assessment:

We received a 3-year, \$225,000(DC) grant from the Arizona Biomedical Research Commission (ABRC) to investigate aging effects in women with ASD. The grant was awarded to Dr. Blair Braden under the New Investigator mechanism. As with the DoD grant, this funding continues the collaboration between Barrow, SARRC, and ASU. The new funding will allow us to include a group of 20 older ASD women and their age-matched TDs. The procedures will mirror our DoD-funded study in order to allow us to directly test gender differences in symptoms, cognition and brain functioning as they age. The funding also allows for longitudinal assessment over 2 time points.

a. What opportunities for training and professional development has the project provided?

As noted in the year 1 annual report, this project provided the environment to train and promote Dr. Baxter's post-doctoral resident, who is one of the key personnel in this study. B. Blair Braden, PhD has been active in all of our projects on ASD and aging. She came to the lab after graduate training in neuroscience, studying hormone and aging in animal models specifically to help develop our cognitive and imaging studies in aging and Autism. She has successfully transitioned to a tenure-track Assistant Professor position at Arizona State University and established a lab in Autism and Aging. Prior to this move when she was still at Barrow Neurological Institute, Dr. Braden applied for a New Investigator Award from the Arizona Biomedical Research Commission. She received her award after she transitioned to her ASU position. Dr. Braden has 1 graduate student and several undergraduate students dedicated to this project. The Barrow lab has one to two Barrett honors students working on this project per year. Through a high school program, the BNI lab had 4 high school students. At the end of the last year, two high school students received 2 INTEL awards for analyses related to this project. These two students are matriculating and attending Harvard University and Washington University in the fall. One of the new students will submit for the Arizona Science Fair this fall.

b. How were the results disseminated to communities of interest?

The following presentations were made or are scheduled:

"Motor System Integrity in Older Adults with Autism Spectrum Disorder."
 Brandon R. Deatherage*, B. Blair Braden, Christopher J. Smith , Michael K. McBeath, Samuel C. McGee , Emily G. Wood, Krishna Sinha, Leslie C. Baxter *Mentored Honors Undergraduate Student (Baxter) International Meeting for Autism Research:, San Francisco, CA May 2017.

- 2) Braden, B.B., Smith, C.J, Thompson, A., Glaspy, T.K., Wood, E., Vatsa, D., and Leslie C. Baxter, L.C. Cognitive and Brain Aging in Autism Spectrum Disorder: Executive Functioning and Frontal and Temporal Lobe Differences. Presented at the International Meeting for Autism Research, San Francisco, CA May 2017, New Adventures in Learning, Chandler-Gilbert Community College, AZ, July 2017.
- 3) Braden, B.B., Riecken, C. Age-Related Cortical Thickness Differences in Adults with Autism Spectrum Disorder. Presented at:; Arizona Alzheimer's Disease Consortium Annual Meeting, May 2017; New Adventures in Learning, Chandler-Gilbert Community College, AZ, July 13, 2017 Society for Neuroscience Annual Meeting, Washington D.C., Nov 2017
- 4) Stoeckmann, M, Baxter, L, Smith, CS, Braden, B*. Large-scale brain networks in middle-aged adults autism spectrum disorder: Functional connectivity differences and relationships with symptoms. Society for Neuroscience, Nov 2017. *Chair for Nanosymposium "Autism: Physiology and Behavior".

Autism Brain Imaging Data Exchange: The Autism Brain Imaging Data Exchange II is large-scale data repository of ASD and TD controls from 17 sites. Our contribution of 58 samples (which includes participants that are part of the DoD study) include the oldest sample in the group (aged 64 years old) and represents the first substantial set of older adults in the exchange. Our contribution to this high-profile, international group of researchers helps us to disseminate our study to the ASD research community.

i. Aging in Autism Special Interest Group, International Society for Autism Research (INSAR).

Both Drs. Baxter and Braden actively participated in the second meeting of the INSAR group of Autism researchers dedicated to the study of older individuals with Autism. The group was formed by Hilde Geurts, PhD of the University of Amsterdam to work towards common goals for studying older adults with ASD. Both Drs. Baxter and Braden are contributing to the establishment of a core set of cognitive tests and other data that can be collated across studies from around the world.

c. What do you plan to do during the next reporting period to accomplish the goals?

- d. Continue with recruitment of ASD and age-matched TDs to reach our target sample of 35/group.
- e. Continue to acquire second time point evaluations of those participants who are two years from their original study.
- f. Begin machine learning data analysis with collaborator Dr. Jieping Ye (University of Michigan) to investigate multimodal influences of aging on ASD.

B) **IMPACT**:

a. What was the impact on the development of the principal discipline(s) of the project?

We published a publication that is one of the first to focus on both behavior and brain imaging in older adults. The reviewers' utilized by the Autism Research editors reflect the importance of our study: "This is an important topic in need of further investigation, and this is a contribution in this direction" and "this work is both timely and important". We have significantly increased the number of older adults that are freely available to others via the national data exchange, "ABIDE II". Our study is complementing the established cognitive studies in older individuals with Autism to emphasize the importance of developing a greater understanding of aging in Autism to inform treatment.

Enhancing studies of the connectome in autism using the autism brain imaging data exchange II. Di Martino, A., O'Connor, D., Chen, B., Alaerts, K, Anderson, J. S., Assaf, M., Balsters, J. H., Baxter, L., Beggiato, A., Bernaerts, S., Blanken, L. M., Bookheimer, S. Y., Braden, B. B., Byrge, L., Castellanos, F. X., Dapretto, M., Delorme, R., Fair, D. A., Fishman, I., Fitzgerald, J., Gallagher, L., Keehn, R. J., Kennedy, D. P., Lainhart, J. E., Luna, B., Mostofsky, S. H., Müller, R. A., Nebel, M. B., Nigg, J. T., O'Hearn, K., Solomon, M., Toro, R., Vaidya, C. J., Wenderoth, N., White, T., Craddock, R. C., Lord, C., Leventhal, B., & Milham, M. P. Sci Data. 2017 March 14;4:170010. doi: 10.1038/sdata.2017.10. PMID: 28291247

What was the impact on other disciplines?

Nothing to Report.

b. What was the impact on technology transfer?

Nothing to Report.

What was the impact on society beyond science and technology?

- i. As reported in the Year 1 summary, an intent of our study is to develop a plan of action to help keep older adults with ASD as independent as possible for as long as possible. We foresee that the results of our study, which is one of the first of its kind, will be able to inform state agencies and community aging programs to develop interventions that will help keep older ASDs independent. We plan on continuing to publish our results, and becoming a voice for the older ASD population, to help form effective and meaningful supports and treatments for this group.
- C) **CHANGES/PROBLEMS:** The Project Director/Principal Investigator (PD/PI) is reminded that the recipient organization is required to obtain prior written approval from the awarding agency Grants Officer whenever there are significant changes in the project or its direction.

If not previously reported in writing, provide the following additional information or state, "Nothing to Report," if applicable:

a. Changes in approach and reasons for change

Nothing to Report

b. Actual or anticipated problems or delays and actions or plans to resolve them

Obtaining institutional and HRPO approval for the study took longer than expected, so recruitment of our cohort is somewhat behind schedule. Our goals are to use all available resources to recruit our target number as soon as possible.

c. Changes that had a significant impact on expenditures

Our expenditures for MRI scans and patient reimbursement is less than expected, as our recruitment has slowed.

d. Significant changes in use or care of human subjects, vertebrate animals, biohazards, and/or select agents

- i. We changed our local consent and HIPAA forms to allow us to share anonymized data with the Autism Brain Imaging Date Exchange (detailed above). This was approved by our institution. We informed the HRPO, who also approved this. Current approval dates are: Informed Consent: 4/20/16; HIPAA: 3/02/16
- e. Significant changes in use or care of human subjects: Nothing to Report
- f. Significant changes in use or care of vertebrate animals. N/A
- g. Significant changes in use of biohazards and/or select agents N/A
- D) **PRODUCTS:** List any products resulting from the project during the reporting period. If there is nothing to report under a particular item, state "Nothing to Report."
- a. **Publications, conference papers, and presentations**Nothing to Report
- i. Journal publications. Nothing to Report
- ii. Books or other non-periodical, one-time publications. Nothing to Report
- b. Other publications, conference papers, and presentations. Nothing to Report

c. Website(s) or other Internet site(s)

Website/link to our media coverage: http://abc7.com/health/adult-men-with-autism-participate-in-one-of-a-kind-study/1429782/

d. List the URL for any Internet site(s) that disseminates the results of the research activities.

A short description of each site should be provided. Nothing to report

e. Technologies or techniques

None to report

f. Inventions, patent applications, and/or licenses

None to report

g. Other Products

None to report

E) PARTICIPANTS & OTHER COLLABORATING ORGANIZATIONS

a. What individuals have worked on the project?

Name:	Leslie C. Baxter, PhD
Project Role:	PI
Researcher Identifier (e.g. ORCID ID):	0000-0002-3971-863X
Nearest person month worked:	<1
Contribution to Project:	Dr. Baxter oversaw data collection, analysis and interpretation and revised a paper submission. She also oversaw data analysis and manuscript prep for 3 papers. Worked with Dr. Braden for grant submissions.
Funding Support:	National Institute on Aging NIA 5 P30 AG019610-03 (Reiman, PI; Baxter, Site PI) State of Arizona, Arizona Alzheimer's Consortium Arizona Biomedical Research Commission

Name:	Brittany Blair Braden, PhD
Project Role:	Co-Investigator
Researcher Identifier (e.g. ORCID ID):	0000-0001-6842-9784
Nearest person month worked:	2
Contribution to Project:	Dr. Braden assisted with oversight of scheduling, data analysis. She is first author of a manuscript that was published. Grant submissions

Funding Support:	Arizona Biomedical Research Commission

- b. Has there been a change in the active other support of the PD/PI(s) or
- c. senior/key personnel since the last reporting period?
- Dr. B. Braden transitioned from post-doctoral fellow to Assistant Professor at Arizona State University. She continues to participate in the study.
- d. What other organizations were involved as partners?
- 1. Organization Name: Southwest Autism Resource and Research Center
- 2. Location of Organization: 2225 N 16th Street Phoenix, AZ 85006
- 3. **Partner's contribution to the project** Partnering PI
- 4. **Collaboration** Recruitment of participants, collaboration with data interpretation and manuscript preparation
- 5. **Organization Name**: Arizona State University
- 6. **Location of Organization:** Department of Speech and Hearing Science, 976 S Forest Mall, Tempe, AZ 85281
- 7. **Partner's contribution to the project** Co investigator
- 8. **Collaboration** Dr. Braden transitioned to an Assistant Professor position at ASU, and her time is now spent overseeing scheduling and data collection, which is done via other funding by Dr. Baxter's research assistants. She focuses on manuscript preparation and grant submissions.